

Thermohaline mixing: A physical mechanism governing the photospheric composition of low-mass giants

Corinne Charbonnel^{1,2} and Jean-Paul Zahn³

¹ Geneva Observatory, University of Geneva, chemin des Maillettes 51, 1290 Sauverny, Switzerland e-mail: Corinne.Charbonnel@obs.unige.ch

² Laboratoire d'Astrophysique de Toulouse et Tarbes, CNRS UMR 5572, Université Paul Sabatier Toulouse 3, 14, av.E.Belin, 31400 Toulouse, France

³ LUTH, CNRS UMR 8102, Observatoire de Paris, 92195 Meudon, France e-mail: Jean-Paul.Zahn@obspm.fr

Accepted for publication in A&A Letter

ABSTRACT

Aims. Numerous spectroscopic observations provide compelling evidence for a non-canonical mixing process that modifies the surface abundances of Li, C and N of low-mass red giants when they reach the bump in the luminosity function. Eggleton et al. (2006) have proposed that a molecular weight inversion created by the $^3\text{He}(^3\text{He},2p)^4\text{He}$ reaction may be at the origin of this mixing, and relate it to the Rayleigh-Taylor instability. In the present work we argue that one is actually dealing with a double diffusive instability referred to as thermohaline convection and we discuss its influence on the red giant branch.

Methods. We compute stellar models of various initial metallicities using the prescription by Ulrich (1972) (extended to the case of a non-perfect gas) for the turbulent diffusivity produced by the thermohaline instability in stellar radiation zones.

Results. Thermohaline mixing simultaneously accounts for the observed behaviour of the carbon isotopic ratio and of the abundances of Li, C and N on the upper part of the red giant branch. It reduces significantly the ^3He production with respect to canonical evolution models as required by measurements of $^3\text{He}/\text{H}$ in galactic HII regions.

Conclusions. Thermohaline mixing is a fundamental physical process that must be included in stellar evolution modeling.

Key words. Instabilities; Stars: abundances, evolution; Galaxies: evolution

1. Introduction

During the first dredge-up (*1dup*; Iben 1965), the surface composition of low-mass red giant stars¹ is modified due to dilution of the external convective stellar layers with hydrogen-processed material: The lithium and the carbon abundances as well as the carbon isotopic ratio decline, while the helium 3 and nitrogen abundances increase. The amplitude of these variations depends on the stellar mass and metallicity. This picture is nicely supported by observations.

After the *1dup* the stellar convective envelope retreats (in mass) ahead of the advancing hydrogen-burning shell (HBS) that surrounds the degenerate helium core, and no further surface abundance variation is predicted by canon-

ical stellar evolution theory² on the RGB. However, spectroscopic observations clearly point out that some non-canonical mixing connects the convective envelope with the HBS and further modifies the surface composition of low-mass giants as soon as they reach the bump in the luminosity function. The sudden drop of the carbon isotopic ratio $^{12}\text{C}/^{13}\text{C}$ provides the most pertinent clue of this mechanism (Gilroy 1989; Gilroy & Brown 1991; Charbonnel 1994; Charbonnel et al. 1998; Gratton et al. 2000; Shetrone 2003; Recio-Blanco & De Laverny 2007) that also modifies the abundances of lithium, carbon and nitrogen (e.g., Gratton et al. 2000). This non-canonical process appears to be universal as it affects at least 95% of the low-mass stars, whether they belong to the field, to open, or globular clusters (Charbonnel & Do Nascimento 1998). This high number satisfies the galactic requirements for the evolution of the ^3He abundance (Tosi 1998; Palla et al. 2000; Romano et al. 2003), since the mecha-

Send offprint requests to: C. Charbonnel

¹ i.e., stars with initial masses below $\sim 2 - 2.5M_{\odot}$ that evolve along the Red Giant Branch (RGB) to high luminosities until helium is ignited in their core under degenerate conditions

² By this we refer to the modelling of non-rotating, non-magnetic stars, where convection is the only transport process considered.

nism which is responsible for the low values of $^{12}\text{C}/^{13}\text{C}$ is also expected to lead to the destruction of ^3He by a large factor in the bulk of the envelope material as initially suggested by Rood et al. (1984; see also Charbonnel 1995; Hogan 1995; Weiss et al. 1996).

At the present time, we have no firm physical model for the non-canonical RGB mixing. Parametrized approaches have been used to reproduce individual observations with the diffusive velocity treated as a free parameter (i.e. Boothroyd & Sackmann 1999; Weiss et al. 2000). On the other hand, rotation-induced mixing has been investigated thoroughly (see references in Charbonnel & Palacios 2004). It turns out however that meridional circulation and shear turbulence alone do not produce enough mixing to account for the surface abundance variations as required by the observations (Palacios et al. 2006).

2. Mixing due to thermohaline convection

Recently Eggleton et al. (2006, 2007) identified a possible cause for such non-canonical mixing, namely the molecular weight inversion created by the $^3\text{He}(^3\text{He}, 2p)^4\text{He}$ reaction in the external wing of the HBS. As it was pointed out already by Ulrich (1972), this nuclear reaction is somewhat singular in that it produces more particles per unit mass than it started from. Using their 3D hydrodynamic code to model a low-mass star at the RGB tip (Dearborn et al. 2006), Eggleton and collaborators found that such a μ -profile leads to efficient mixing, of the kind required to “reconcile Big Bang and stellar nucleosynthesis (as far as ^3He is concerned)”, as they put it.

According to them, the instability responsible for that mixing is the well-known Rayleigh-Taylor instability, a dynamical instability which is triggered when a layer of heavier fluid lies over lighter fluid. In stellar interiors, that instability takes the form of the convective instability, which tends to render the temperature gradient adiabatic rather than to suppress the density inversion. But what first occurs in a star, as the inverse μ -gradient builds up, is actually a double diffusive instability, which was discussed long ago in the literature under the generic name of “thermohaline convection” (Stern 1960). This instability appears in various astrophysical situations, for instance when ^4He or C-rich material is deposited at the surface of a star in a mass transferring binary (Stothers & Simon 1969; Stancliffe et al. 2007). Recently it has been invoked when a star accretes heavy elements during planet formation (Vauclair 2004).

The thermohaline instability differs from the convective instability in that it involves two components, of which one, the stabilizing one (temperature) diffuses faster than the other (salt), whose stratification is unstable. It occurs in a stable stratification that satisfies the Ledoux criterion for convective stability:

$$\nabla_{\text{ad}} - \nabla + \left(\frac{\varphi}{\delta}\right) \nabla_{\mu} > 0, \quad (1)$$

but where the molecular weight decreases with depth:

$$\nabla_{\mu} := \frac{d \ln \mu}{d \ln P} < 0. \quad (2)$$

We use the classical notations for $\nabla = (\partial \ln T / \partial \ln P)$, $\varphi = (\partial \ln \rho / \partial \ln \mu)_{P,T}$ and $\delta = -(\partial \ln \rho / \partial \ln T)_{P,\mu}$.

In the laboratory, the instability takes the form of “salt fingers”; since heat diffuses faster than salt, these fingers sink because they grow increasingly heavier than their environment, until they become turbulent and dissolve. In stellar interiors, the role of salt is played by a heavier species, such as helium, in a hydrogen-rich medium.

Ulrich (1972) was the first to derive a prescription for the turbulent diffusivity produced by that instability in stellar radiation zones. Through a linear analysis, and assuming perfect gas ($\varphi = \delta = 1$) he gets (his eqs. 24 and 31)

$$D_t = C_t K \frac{-\nabla_{\mu}}{(\nabla_{\text{ad}} - \nabla)} \quad \text{for } \nabla_{\mu} < 0; \quad (3)$$

K is the thermal diffusivity. His non-dimensional coefficient involves the aspect ratio α (length/width) of the fingers:

$$C_t = \frac{8}{3} \pi^2 \alpha^2; \quad (4)$$

for the value he advocates, $\alpha = 5$, this coefficient is rather large: $C_t = 658$.

Kippenhahn et al. (1980) extended Ulrich’s expression to the case of a non-perfect gas (including radiation pressure, degeneracy):

$$D_t = C_t K \left(\frac{\varphi}{\delta}\right) \frac{-\nabla_{\mu}}{(\nabla_{\text{ad}} - \nabla)} \quad \text{for } \nabla_{\mu} < 0. \quad (5)$$

However instead of fingers they considered “blobs” which according to them are destroyed very rapidly: their mixing length is of the order of the blob size and this translates into a smaller value of 12 for the coefficient C_t .

It is of course disturbing that these coefficients C_t differ by almost two orders of magnitude, depending on whom one believes; it would be most desirable to confront them with realistic 3D simulations. In the meanwhile, we tend to favour a large value for the coefficient C_t as advocated by Ulrich, because all experiments so far have displayed slender fingers, rather than blobs (cf. Krishnamurti 2003). Crude as it may be, this prescription has the advantage of being rooted in the physical process, unlike the admittedly ad hoc diffusivity chosen by Eggleton et al. (2007).

3. Model calculations

We compute three evolution models of a $0.9 M_{\odot}$ star with various initial values of $[\text{Fe}/\text{H}]$, namely -1.8 , -1.3 and -0.5 , with the aim of comparing our theoretical predictions with observations over a large range of metallicity. We use the code STAREVOL (Siess et al. 2000; Palacios et al. 2003, 2006). In the present study the transport of particles in the radiative regions is due to thermohaline mixing only.




Fig. 1. Profiles of the logarithm of the mass fraction of H, ^3He , ^{12}C , ^{13}C , ^{14}N , ^{16}O , and of the mean molecular weight gradient $\nabla_\mu = d \ln \mu / d \ln P$ (bold line, full when $\nabla_\mu > 0$, dashed otherwise) as a function of the reduced mass (see text) inside a $0.9 M_\odot$, $[\text{Fe}/\text{H}] = -1.3$ star. The left and right panels correspond to models including thermohaline mixing and located just before and after the bump. In the latter case, the thermohaline instability has just set in naturally, while in the former case it is still prevented due to the strong μ -gradient left behind by the *1dup*

The expression we use for D_t is that given in Eq. 5 that includes the correction for a non-perfect gas. For simplicity we assume $C_t = 1000$, which is of the order of magnitude of the coefficient advocated by Ulrich (1972). The model at $[\text{Fe}/\text{H}] = -1.3$ is also computed with $C_t = 100$ and 12. We do not consider any additional transport process related to rotation.

Fig. 1 presents the abundance profiles of selected elements and the mean molecular weight gradient $\nabla_\mu = d \ln \mu / d \ln P$, just before and after the bump in the $0.9 M_\odot$, $[\text{Fe}/\text{H}] = -1.3$ star when $C_t = 1000$. The abscissa is the relative mass $\delta M = (M_r - M_{\text{core}}) / (M_{\text{env}} - M_{\text{core}})$ defined as ranging from 0 to 1 between the bottom of the HBS and the base of the convective envelope.

During the *1dup* phase, the convective envelope homogenizes the star down to very deep regions, and builds a very steep gradient of molecular weight at the point of its maximum penetration. This corresponds to the external peak (at $\delta M \sim 0.12$) in the profile of ∇_μ that can be seen in the left panel of Fig. 1. On the other hand the deeper peak (at $\delta M \sim 0.05$) locates the region where H is efficiently depleted by nuclear reactions in the HBS. One sees that at the current luminosity ($L = 90 L_\odot$) both peaks are very close to each other, meaning that the star is approaching the bump. At this evolutionary point, ∇_μ




Fig. 2. Logarithm of the diffusion coefficient D_t (assuming $C_t = 1000$) inside the $0.9 M_\odot$, $[\text{Fe}/\text{H}] = -1.3$ star just after the bump.

is positive in the whole radiative region, and the thermohaline instability cannot set in (see Eq. 2). The profiles of the chemical elements are thus identical to those obtained in a canonical model.

When the HBS passes through the μ -discontinuity left behind by the *1dup* (i.e., at the bump), H-burning occurs in a homogeneous region. The $^3\text{He}(^3\text{He}, 2p)^4\text{He}$ reaction slightly lowers the molecular weight in the external wing of the HBS where ∇_μ becomes negative (bold dashed line in the right panel of Fig. 1), allowing the thermohaline instability to develop naturally between the ^3He -burning region and the base of the convective envelope. Deeper inside the radiative region, ∇_μ remains positive (bold full line) and no thermohaline mixing occurs. As can be seen in Fig. 1, the surface abundance of ^3He , ^{12}C , ^{13}C , and ^{14}N are already modified soon after the onset of thermohaline mixing. However the surface abundance of ^{16}O remains constant because the thermohaline mixing does not extend down to the very deep region where full CNO-burning operates at equilibrium.

We show in Fig. 2 the profile of the diffusion coefficient D_t (Eq. 5, for $C_t = 1000$), for the same model just after the onset of the thermohaline instability ($L = 115 L_\odot$). D_t is high enough to connect the HBS and the convective envelope and to modify the surface abundances as shown below. It is two to three orders of magnitude higher than that characterizing the rotational mixing (Palacios et al. 2006).

4. Signatures of mixing and comparison with observations

In Fig. 3 we compare the temporal evolution of ^7Li , $^{12}\text{C}/^{13}\text{C}$, $[\text{C}/\text{Fe}]$, and $[\text{N}/\text{Fe}]$, obtained at the surface of our models with homogeneous observational data for field stars with $-2 \leq [\text{Fe}/\text{H}] \leq -1$ by Gratton et al. (2000). We see




Fig. 3. Evolution of the lithium abundance, of the carbon isotopic ratio $^{12}\text{C}/^{13}\text{C}$, of $[\text{C}/\text{Fe}]$ and $[\text{N}/\text{Fe}]$ as a function of the luminosity logarithm for the $0.9M_{\odot}$ models with $[\text{Fe}/\text{H}]=-1.8$, -1.3 and -0.5 (respectively dashed-dotted blue, full line black, and dashed red) computed with $C_t=1000$ in Eq. 5. The black dotted line is for the model with $[\text{Fe}/\text{H}]=-1.3$ calculated with $C_t=100$. The arrows in the upper panel indicate the location of the bump for the three metallicities. Observational data are from Gratton et al. (2000) for field stars in the metallicity range $[\text{Fe}/\text{H}] \in [-2; -1]$. Blue symbols are for stars with measured $[\text{Fe}/\text{H}]$ values lower than -1.4 , black for stars with $-1.4 \leq [\text{Fe}/\text{H}] \leq -1.2$, and red for $[\text{Fe}/\text{H}]$ higher than -1.2 . Circles are actual measurements, open upward triangles are *lower* limits and open downward triangles are *upper* limits.

first that variations associated with the *1dup* fit well the observational behaviour on the lower RGB. The surface abundances then stay constant until the stars reach the bump. Until that evolutionary point the predictions are identical to those of the canonical models.

Then at the luminosity of the bump (which increases for a given stellar mass when the initial metallicity decreases, as shown by the arrows) thermohaline mixing leads to a second episode of abundance variations that explains remarkably well the data³ Li is rapidly destroyed, and the carbon isotopic ratio reaches values between 5 and 7 very close from the equilibrium value. $[\text{C}/\text{Fe}]$ and

³ Oscillations in stellar luminosity can be seen just above the bump. They are related to the transport of ^3He from the convective envelope into the outer HBS region where this element burns in favour of ^7Be . The related nuclear energy release is large. When it happens to rival that of the pp-chains, the total luminosity increases and the star readjusts. Once the abundance of ^3He is low enough the oscillations vanish.

$[\text{N}/\text{Fe}]$ respectively decrease and increase. These C and N variations are stronger for lower initial metallicity, as required by the observations. As already noted in § 3, the surface abundances of the heavier elements (^{16}O and ^{23}Na in particular, not shown here) which are affected by nuclear reactions much closer to the He-core do not vary at the surface.

The models shown in Fig. 3 were computed assuming $C_t = 1000$ in Eq. 5. We also show the predictions for the model with $[\text{Fe}/\text{H}]=-1.3$ computed with $C_t = 100$. In that case the surface abundances change less rapidly and one does not reproduce the abrupt variation at the luminosity of the bump depicted by the observations. Our analysis is thus compatible with the high value for the coefficient C_t advocated by Ulrich (1972) and it excludes the low value (12, not shown here) derived by Kippenhahn et al. (1980).

5. The case of ^3He

On the main sequence, a ^3He peak builds up due to pp-reactions inside the low-mass stars (Iben 1967), and is engulfed in the stellar envelope during the *1dup*. As a consequence the surface abundance of ^3He strongly increases on the lower RGB as can be seen in Fig. 4 for our models at three different metallicities. Its value reaches a maximum when the whole peak is engulfed. After the *1dup* the temperature at the base of the convective envelope is too low for ^3He to be nuclearly processed. As a result in canonical models ^3He stays constant at the surface and its final value is strongly increased with respect to the initial one (this is the value before thermohaline mixing sets in at the bump).

After the bump, thermohaline mixing brings ^3He from the convective envelope down to the region where it is nuclearly burned. This leads to a rapid decreases of the surface abundance of this element as can be seen in Fig. 4. This confirms the early suggestion by Rood et al. (1984) that the variations of the carbon isotopic ratio and of ^3He are strongly connected (see also Charbonnel 1995; Eggleton et al. 2007). It is important to note that in the models presented here ^3He decreases by a large factor in the ejected material with respect to the canonical evolution predictions but that low-mass stars remain net producers (while far much less efficient than in the canonical case) of ^3He . Computations for different masses and metallicities have now to be performed in order to estimate the actual contribution of low-mass stars to Galactic ^3He in the framework proposed here. We are confident that the corresponding ^3He yields will help reconciling the primordial nucleosynthesis with measurements of $^3\text{He}/\text{H}$ in galactic HII regions (Balser et al. 1994; Bania et al. 2002; Charbonnel 2002).

6. Conclusions

It is known since a long time that a non-canonical mixing process modifies the surface abundances of low-mass stars when they arrive at the bump in the luminosity function.




Fig. 4. Evolution of the surface abundance of ^3He (in mass fraction). The line symbols are as in Fig. 3.

Based on 3D-modeling of a low-mass star at the RGB tip, Eggleton and collaborators have proposed that the molecular inversion created by the $^3\text{He}(^3\text{He},2p)^4\text{He}$ reaction in the external wing of the HBS may be the root cause for such a mixing. They ascribe this mixing to the Rayleigh-Taylor instability.

In the present work we argue that one is actually dealing with a double diffusive instability called thermohaline convection, which has been discussed long ago in the literature. We compute stellar models including the prescription given by Ulrich (1972) and extended to the case of a non-perfect gas for the turbulent diffusivity produced by that instability in stellar radiative zone. The results presented here indicate that thermohaline convection simultaneously accounts for the observed behaviour of the carbon isotopic ratio and for the abundances of Li, C and N in RGB stars at and above the bump in the luminosity function. It also avoids large ^3He production by low-mass stars as required by chemical evolution models of the Galaxy. It does not modify the O nor the Na surface abundances. In a future work the effect of the thermohaline mixing in models of various masses will be presented.

Acknowledgements. We thank the referee for her/his comments that helped us improve our manuscript and Dr Eggleton for sending us his paper before publication. C.C. is supported by the Swiss National Science Foundation (FNS). We acknowledge travel support from the French Programme National de Physique Stellaire (PNPS).

References

- Balser, D.A., Bania, T.M., Brockway, C.J., Rood, R.T., Wilson, T.L., 1994, *ApJ*, 430, 667
 Bania, T.M., Rood, R.T., Balser, D.A., 2002, *Nature*, 415, 54
 Boothroyd, A.I., Sackmann, I.J., 1999, *ApJ*, 510, 232
 Charbonnel, C., 1994, *A&A* 282,811

- Charbonnel, C., 1995, *ApJ Letters*, 453, 41
 Charbonnel, C., 2002, *Nature*, 415, 27
 Charbonnel, C., Brown, J.A., Wallerstein, G., 1998, *A&A* 332, 204
 Charbonnel, C., 2005, IAU Symp.228 on “From lithium to uranium : Element tracers of early cosmic evolution”, Cambridge Univ.Press, Eds. V. Hill, P. François, F. Primas, p.347
 Charbonnel, C., do Nascimento, J.D., 1998, *A&A* 336, 915
 Charbonnel, C., Palacios, A., 2004, IAU Symp.215, 440
 Dearborn, D.S.P., Eggleton, P.P., Lattanzio, J.C., 2006, *ApJ*, 639, 405
 Eggleton, P.P., Dearborn, D.S.P., Lattanzio, J.C., 2006, *Science*, 314, 1580
 Eggleton, P.P., Dearborn, D.S.P., Lattanzio, J.C., 2007, private communication
 Gilroy, K.K., 1989, *ApJ* 347, 835
 Gilroy, K.K., Brown, J., 1991, *ApJ* 371, 578
 Gratton, R.G., Sneden, C., Carretta, E., Bragaglia, A., 2000, *A&A* 354, 169
 Hogan, C.J., 1995, *ApJ Letters*, 441, 17
 Iben, I., 1967, *ApJ*, 143, 642
 Kippenhahn, R., Ruschenplatt, G., Thomas, H.C., 1980, *A&A*, 91, 175
 Krishnamurti, R., 2003, *J. Fluid Mech.*, 483, 287
 Palacios, A., Talon, S., Charbonnel, C., Forestini, M., 2003, *A&A* 399, 603
 Palacios, A., Charbonnel, C., Talon, S., Siess, L., 2006, *A&A*, 453, 261
 Palla, F., Bachiller, R., Stanghellini, L., Tosi, M., Galli, D., 2000, *A&A*, 355, 69
 Recio-Blanco, A., de Laverny, P., 2007, *A&A*, 461, L13
 Romano, D., Tosi, M., Matteucci, F., Chiappini, C., 2003, *MNRAS*, 346, 295
 Rood, R.T., Bania, T.W., Wilson, T.L., 1984, *ApJ*, 280, 629
 Shetrone, M.D., 2003, in *CNO in the Universe*, eds Charbonnel C., Schaerer D., Meynet G., ASP Conference Series 304, 137
 Siess, L., Dufour, E., Forestini, M., 2000, *A&A* 358, 593
 Stancliffe, R.J., Glebbeek, E., Izzard, R.G., Pols, O.R., 2007, *AALetter* in press
 Stern, M.E., 1960, *Tellus*, 12, 172
 Stothers, R., Simon, N.R., 1969, *ApJ*, 157, 673
 Tosi, M., 1998, *Space Science Reviews*, Vol.84, 207
 Ulrich, R.K., 1972, *ApJ*, 172, 165
 Vauclair, S., 2004, *ApJ*, 605, 874
 Weiss, A., Wagenhuber, J., Denissenkov, P.A., 1996, *A&A*, 313, 581
 Weiss, A., Denissenkov, P.A., Charbonnel, C., 2000, *A&A*, 356, 181